

Policy

# Clinical Guideline

## South Australian Perinatal Practice Guidelines – Neonatal Hypoglycaemia

**Policy developed by:** SA Maternal and Neonatal Clinical Network

**Approved SA Health Safety & Quality Strategic Governance Committee on:**

8 October 2013

**Next review due:** 30 August 2016

**Summary** Guideline for the management of the neonate with hypoglycaemia

**Keywords** morbidity, hypoglycaemia, jitteriness, ketogenesis, caesarean, diabetes, hypothermia, breast feed, seizures, respiratory, erythroblastosis fetalis, prematurity, Perinatal Practice Guidelines, Neonatal Hypoglycaemia, clinical guideline

**Policy history** Is this a new policy? **N**  
Does this policy amend or update an existing policy? **Y**  
Does this policy replace an existing policy? **Y**  
If so, which policies? Neonatal Hypoglycaemia

**Applies to** All SA Health Portfolio  
All Department for Health and Ageing Divisions  
All Health Networks  
CALHN, SALHN, NALHN, CHSALHN, WCHN, SAAS  
Other

**Staff impact** All Clinical, Medical, Nursing, Allied Health, Emergency, Dental, Mental Health, Pathology

**PDS reference** CG116

---

### Version control and change history

Version	Date from	Date to	Amendment
1.0	18/02/2008	01/04/2008	Original version
2.0	01/04/2008	18/01/2011	Reviewed
3.0	18/01/2011	23/09/2013	Reviewed
4.0	23/09/2013	Current	



# South Australian Perinatal Practice Guidelines

# neonatal hypoglycaemia

© Department of Health, Government of South Australia. All rights reserved.

## Note

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

**ISBN number:**  
**Endorsed by:**  
**Last Revised:**  
**Contact:**

978-1-74243-084-3  
South Australian Maternal & Neonatal Clinical Network  
23/09/13  
South Australian Perinatal Practice Guidelines Workgroup at:  
cywhs.perinatalprotocol@health.sa.gov.au




Government  
of South Australia  
SA Health

# South Australian Perinatal Practice Guidelines neonatal hypoglycaemia

© Department of Health, Government of South Australia. All rights reserved.

## Management of hypoglycaemia in the newly born and neonatal periods



## Important points

### Key physiological principles<sup>1,2,3</sup>

- > In the newly born, glucose levels are maintained in blood by the release of liver glycogen reserves in the first few hours, by gluconeogenesis, and by sparing of glucose consumption through the oxidation of fat reserves and the utilisation of ketone bodies and lactic acid as energy substrates by the brain.
- > Ketogenesis is the adaptive response of principal importance. Utilisation of ketones by the brain allows normal cerebral energy metabolism at lower plasma glucose levels in babies in comparison to older children and adults.
- > Newly born babies with impaired adaptive ketogenesis are considered 'at risk' of impaired cerebral energy metabolism at low levels of plasma glucose. This includes particularly infants who are preterm, growth restricted, born to mothers with gestational or insulin dependent diabetes, or who have an abnormal transition period due to adverse perinatal circumstances.

### Measurement of plasma glucose is required in two broad groups of babies

- > Newly born, 'at risk' babies who are otherwise well. These babies potentially have impaired adaptive ketogenesis and increased glucose utilisation. They are screened for hypoglycaemia with plasma glucose levels at 1 and 4 hours of age to capture the nadir of plasma glucose.<sup>4</sup>
- > Babies who are identified as unwell, who present with neonatal seizures, or who are receiving normal postnatal care and develop symptoms consistent with hypoglycaemia and impaired cerebral energy metabolism. Plasma glucose is measured in these babies immediately at presentation.

### Principles and pitfalls in measurement of plasma glucose

- > Wide variations in plasma glucose results may occur due to errors in sample collection, processing and assay method. This is particularly important to consider in small hospitals with limited laboratory support.
- > Glycolysis by red cells is rapid in newborn samples. Blood samples sent to the laboratories for glucose estimation need to be collected into fluoride containers (preferably), or centrifuged immediately to separate red cells, or kept on ice to avoid inaccuracies due to red cell metabolism.
- > Glucose analysis using standard laboratory methods, blood gas machines or ISTAT devices is accurate provided the sample is processed immediately.
- > Glucometer and HemaCue machines have utility because they are simple and give immediate bed-side results. However, they may have serious inaccuracies at levels of plasma glucose around commonly applied neonatal intervention thresholds. The main danger is a false negative reading where a 'normal' result is given while the actual plasma glucose is abnormally low. Therefore, if using these machines, this guideline recommends adjusting the intervention threshold upward to avoid false negatives.

### Understanding the definition of hypoglycaemia and how plasma glucose levels are used to guide management

**ISBN number:** 978-1-74243-084-3  
**Endorsed by:** South Australian Maternal & Neonatal Clinical Network  
**Last Revised:** 23/09/13  
**Contact:** South Australian Perinatal Practice Guidelines Workgroup at:  
cywhs.perinatalprotocol@health.sa.gov.au

- > Hypoglycaemia is defined physiologically as that blood glucose level at which cerebral energy needs fail to be met.
- > However, there is no clear and satisfactory definition of hypoglycaemia based on plasma glucose levels alone, because of the complexities of metabolic and hormonal adaptation to birth, our limited understanding of neonatal cerebral defences in hypoglycaemia, and inadequate long-term data to fully assess the impact of hypoglycaemia on the developing brain<sup>1,5,6</sup>
- > Therefore, 'Operational' or 'Intervention' thresholds are used, which are pragmatic treatment levels for plasma glucose that aim to balance safety with the avoidance of over treatment of healthy babies<sup>6</sup>
- > Once a decision has been made to treat hypoglycaemia, a 'Therapeutic' goal is set that is deliberately higher than the threshold for intervention to allow a safety margin for management
- > The intervention threshold is lower for the newly born 'at risk' baby than for older or symptomatic babies because plasma glucose normally falls after birth and there is a desire to avoid over-treating normal babies and interfering with breast feeding

## Summary of plasma glucose intervention thresholds and therapeutic goals<sup>6</sup>

Quoted values are for gold standard laboratory methodology

Clinical setting	Intervention threshold	Therapeutic goal	Main therapeutic intervention
Well infant, not in 'at risk' group	Don't measure glucose unless hypothermic, feeding abnormally, or becomes unwell.		Normal care
'At risk' well infant screened for hypoglycaemia at 1 and 4 hours of age	< 2.0 mmol / L	> 2.5 mmol / L	Feeding
Unwell infant Abnormal feeding or hypothermia beyond the 4 hour transition period Seizures Requiring intravenous fluids (implies an unwell infant)	< 2.5 mmol / L	> 2.5-3.5 mmol / L	IV 10 % glucose. Feeds in selected cases.
All infants	< 1.5 mmol / L urgent treatment required with IV 10 % glucose	> 3.5 mmol / L	IV 10 % glucose

### Important points of emphasis are:

- > Healthy term babies who are not 'at risk' should not have plasma glucose levels measured because the result will have little meaning<sup>4,6</sup>
- > Plasma glucose levels of < 1.5 mmol / L require urgent treatment with IV 10 % glucose to avoid cerebral energy deficiency and potential brain injury
- > In utero growth restriction is an important cause of hypoglycaemia and other morbidity. For babies who are significantly clinically malnourished (little subcutaneous fat, loose skin folds, withered appearance) formula supplementation of breast-feeding until breast milk flow is established is recommended (see PPG: Fetal growth restricted)
- > Disinterest in, or poor feeding for more than 8 hours after birth, or after previous feed, (particularly when previous feeding has been good) requires medical assessment and a plasma glucose measurement
- > All babies born to mothers with diabetes are at risk of hypoglycaemia regardless of size at birth or the tightness of glycaemic control in pregnancy, and all such babies should be screened with plasma glucose levels.<sup>1</sup> Mothers with diabetes should be encouraged to express colostrum / breast milk and store frozen for later use

**ISBN number:** 978-1-74243-084-3  
**Endorsed by:** South Australian Maternal & Neonatal Clinical Network  
**Last Revised:** 23/09/13  
**Contact:** South Australian Perinatal Practice Guidelines Workgroup at:  
 cywhs.perinatalprotocol@health.sa.gov.au

- > Large size for gestation (see PPG: accelerated fetal growth) is not a risk factor for hypoglycaemia where glucose challenge or tolerance tests have been documented to be normal, and such babies do not require routine screening<sup>7,8</sup>
- > Jitteriness is common and usually benign, and a PGL > 2.5 mmol / L excludes hypoglycaemia as a cause
- > Where a baby is 'at risk' or has symptoms consistent with hypoglycaemia and a plasma glucose level can't be obtained or the result will be delayed, appropriate treatment with feeding or IV fluids should be instituted and not deferred until the glucose result is known
- > **In most cases of documented hypoglycaemia, paediatric or neonatology advice should be sought due to the many considerations in management both of the hypoglycaemia and associated conditions**
- > **Unusual or persistent hypoglycaemia will require further investigation for metabolic or endocrine causes, and such babies should not be discharged without further evaluation**

## Guidelines for screening, detection and management of hypoglycaemia in the newly born and neonatal periods

### 1. Newly born who are well but identified as 'at risk'

Well, transitional babies are at risk of hypoglycaemia if there are perinatal factors that impair adaptive ketogenesis and/or increase glucose consumption

- > Respiratory distress post elective caesarean birth at term with a low level oxygen requirement ( $\text{FIO}_2 < 0.3$ ) and suspected transient tachypnoea of the newborn
- > Maternal diabetes
- > Birth weight  $< 10^{\text{th}}$  percentile for gestation
- > Mild prematurity (32-37 weeks gestation)
- > Hypothermia (per rectum temperature  $< 36^\circ\text{C}$ )
- > Suspected perinatal asphyxia where there has been a need for simple resuscitation measures (includes positive pressure with face mask where there is a rapid response to resuscitation but a need for transitional observation)
- > Maternal treatment with drugs that interfere with glucose homeostasis, most importantly beta-blockers and valproic acid

These babies have screening plasma glucose levels at 1 and 4 hours of age

If there is no respiratory distress the baby should be offered the breast as soon as practicable, or if the mother wishes to artificially feed offered a bottle of formula

If there is low level respiratory distress, feeding is not required in the first 4 hours provided PGLs remain normal

#### ***If both the 1 hour and 4 hour PGL $\geq 2.0$ mmol / L (normal)***

- > These babies should be treated as normally as possible
- > Seek advice if further clinical concerns

#### ***If either the 1 hour or 4 hour PGL is 1.5 - 2.0 mmol / L***

- > If no respiratory distress, these babies should be fed
- > Breast feed if mother available, or if mother wishes to breast feed but is unavailable, give EBM or 10 % glucose orally in the quantity equivalent to a one or two hourly aliquot of a standard starting volume (around 2-4 mL/kg) by a means agreeable to mother and staff. A similar formula volume can be given with mother's consent or if her intent is to artificially feed
- > If respiratory distress, give intravenous 10 % glucose bolus (2 mL / kg) over 5 minutes followed by an infusion of 60 mL/kg/day of 10 % glucose
- > Repeat PGL in 30 to 60 minutes. Therapeutic goal PGL  $> 2.5$ mmol / L for enteral feeding and  $> 3.5$  mmol / L if on IV fluids
- > Seek advice

#### ***If PGL $< 1.5$ mmol / L at any time***

- > Obtain IV access and give 10 % glucose bolus (as above) and 10 % glucose infusion initially at 90 mL/kg/day above. Consider glucagon 0.1 milligram / kg (= 0.1 units / kg) IM, if IV access delayed

**ISBN number:** 978-1-74243-084-3  
**Endorsed by:** South Australian Maternal & Neonatal Clinical Network  
**Last Revised:** 23/09/13  
**Contact:** South Australian Perinatal Practice Guidelines Workgroup at:  
cywhs.perinatalprotocol@health.sa.gov.au

- > Repeat PGL in 30 minutes
- > Therapeutic goal PGL > 3.5 mmol / L. Consider adjusting 10 % glucose infusion rate down if threshold achieved
- > Seek advice

## 2. Postnatal ward babies

A PGL and medical assessment is immediately advised for postnatal ward babies with:

- > Disinterest in feeding for 8 hours or more from birth or the last feed
- > Hypothermia (PR temp < 36°C)
- > Jitteriness

### **If PGL ≥ 2.5 mmol / L**

- > Hypoglycaemia is not the cause of symptoms

### **If PGL 1.5-2.5 mmol / L**

- > Feeding is attempted with a repeat PGL in 30-60 minutes
- > Therapeutic goal PGL > 2.5 mmol / L; consider 10 % glucose infusion at 60 mL/kg/day if goal not achieved
- > Seek advice

### **If PGL < 1.5 mmol / L**

- > Obtain IV access and give an IV 10 % glucose bolus (as above) and 10 % glucose infusion initially at 90 mL/kg/day. Consider glucagon 0.1 milligram / kg (= 0.1 units / kg) IM, if IV access delayed
- > Repeat PGL in 30 minutes.
- > Therapeutic goal PGL > 3.5 mmol / L. Consider adjusting 10 % glucose infusion rate down if threshold achieved
- > Seek advice

## 3. Babies at any age with a major illness

Immediate IV access and plasma glucose measurement are performed at any age in specific clinical contexts where an illness has become apparent since birth or there are neurological symptoms consistent with impaired neuronal metabolism

- > Seizures
- > Unwell baby
  - > Ongoing or evolving respiratory illness
  - > Suspected or proven infection with clinical instability
  - > Suspected congenital heart disease presenting with cyanosis or poor perfusion
  - > Suspected perinatal asphyxia where there has been a need for intubation or cardiac massage during resuscitation
  - > Erythroblastosis fetalis
- > Significant prematurity (< 32 weeks gestation)

All such babies will require an intravenous 10 % glucose infusion commencing at 60 mL/kg/day as part of management of the underlying problem

A PGL is taken immediately at the time of assessment

**ISBN number:** 978-1-74243-084-3  
**Endorsed by:** South Australian Maternal & Neonatal Clinical Network  
**Last Revised:** 23/09/13  
**Contact:** South Australian Perinatal Practice Guidelines Workgroup at:  
cywhs.perinatalprotocol@health.sa.gov.au

***If PGL < 2.5 mmol / L***

- > Give an IV 10 % glucose bolus 2 mL / kg (as above) and 10 % glucose infusion initially at 90 mL/kg/day. Consider glucagon 0.1 milligram / kg (= 0.1 units / kg) IM, if IV access delayed
- > Repeat the PGL in 30 minutes
- > Therapeutic goal PGL > 3.5 mmol / L. Consider adjusting 10 % glucose infusion rate down if threshold achieved
- > Ongoing paediatric or neonatology advice is required

**ISBN number:**  
**Endorsed by:**  
**Last Revised:**  
**Contact:**

978-1-74243-084-3  
South Australian Maternal & Neonatal Clinical Network  
23/09/13  
South Australian Perinatal Practice Guidelines Workgroup at:  
cywhs.perinatalprotocol@health.sa.gov.au

## References

1. Platt MW, Deshpande S. Metabolic adaptation at birth. *Sem Fetal Neonatal Med* 2005;10:341-350
2. Hawdon JM et al. Patterns of metabolic adaptation for preterm and term infants in the first neonatal week. *Arch Dis Child* 1992;67:357-365
3. Hawdon JM, Ward Platt MP. Metabolic adaptation in small for gestational age infants. *Arch Dis Child* 1993;68:262-268
4. Deshpande S, Ward Platt M. The investigation and management of neonatal hypoglycaemia. *Sem Fet Neonat Med* 2005;10:351-361
5. Boluyt N et al. Neurodevelopment after neonatal hypoglycaemia: a systematic review and design of an optimal future study. *Pediatrics* 2006;117(6):2231-2243
6. Cornblath M et al. Controversies regarding definition of neonatal hypoglycaemia: suggested operational thresholds. *Pediatrics* 2000; 105:1141-1145
7. Lubchenco LO, Bard H. Incidence of hypoglycaemia in newborn infants classified by birth weight and gestational age. *Pediatrics* 1971; 47(5): 831-838
8. Brand PLP, Molenaar NLD, Kaaijk C, Wierenga WS. Neurodevelopmental outcome of hypoglycaemia in healthy, large for gestational age, term newborns. *Arch Dis Child* 2005;90(1):78-81

## Abbreviations

BW	Birth weight
C	Celsius
EBM	Expressed breast milk
et al	And others
IM	Intramuscular
IUGR	Intrauterine growth restriction
IV	Intravenous
kg	Kilogram(s)
mL	Millilitre(s)
mmol/L	Millimols per litre
PR	Per rectum
PGL	Plasma glucose level
Prem	Premature
TTN	Transient tachypnoea of the newborn