

Perinatal Practice Guideline

Clinical Guideline

Sepsis in Pregnancy

Policy developed by: SA Maternal, Neonatal & Gynaecology Community of Practice

Approved SA Health Safety & Quality Strategic Governance Committee on: 01 March 2017

Next review due: 31 March 2020

Summary The purpose of this guideline is to provide clinicians with information and treatment guidelines for sepsis in pregnancy. It particularly focuses on recognition of risk factors, diagnosis and management of sepsis in the peri partum period, including Early Goal Directed Therapy in the case of severe sepsis.

Keywords Sepsis in Pregnancy, sepsis, chorioamnionitis, infection, antibiotics, toxic shock syndrome, clinical guideline, PPG, perinatal practice guideline, early goal directed therapy, toxic shock syndrome, endometritis, bacteraemia, systemic inflammatory response syndrome, septicaemia

Policy history Is this a new policy? **N**
Does this policy amend or update an existing policy? **Y v1.0**
Does this policy replace an existing policy? **Y**
If so, which policies?

Applies to All Health Networks
CALHN, SALHN, NALHN, CHSALHN, WCHN, SAAS

Staff impact All Staff, Management, , Students, All Clinical, Medical, Midwifery, Nursing, Allied Health, Emergency, Mental Health, Pathology

PDS reference CG190

Version control and change history

Version	Date from	Date to	Amendment
1.0	20 Nov 12	28 Feb 17	Original version
2.0	28 Feb 17	6 March 2017	Reviewed
3.0	01 March 2017	Current	Review and update

© Department for Health and Ageing, Government of South Australia. All rights reserved.

South Australian Perinatal Practice Guidelines

Sepsis in Pregnancy

© Department of Health, Government of South Australia. All rights reserved.

Note

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

Explanation of the Aboriginal artwork:

The Aboriginal artwork used symbolises the connection to country and the circle shape shows the strong relationships amongst families and the Aboriginal culture. The horse shoe shape design shown in front of the generic statement symbolises a woman and those enclosing a smaller horse shoe shape depicts a pregnant women. The smaller horse shoe shape in this instance represents the unborn child. The artwork shown before the specific statements within the document symbolises a footprint and demonstrates the need to move forward together in unison.



Australian Aboriginal Culture is the oldest living culture in the world yet Aboriginal people continue to experience the poorest health outcomes when compared to non-Aboriginal Australians. In South Australia, Aboriginal women are 2-5 times more likely to die in childbirth and their babies are 2-3 times more likely to be of low birth weight. The accumulative effects of stress, low socio economic status, exposure to violence, historical trauma, culturally unsafe and discriminatory health services and health systems are all major contributors to the disparities in Aboriginal maternal and birthing outcomes. Despite these unacceptable statistics the birth of an Aboriginal baby is a celebration of life and an important cultural event bringing family together in celebration, obligation and responsibility. The diversity between Aboriginal cultures, language and practices differ greatly and so it is imperative that perinatal services prepare to respectively manage Aboriginal protocol and provide a culturally positive health care experience for Aboriginal people to ensure the best maternal, neonatal and child health outcomes.

Purpose and Scope of PPG

The purpose of this guideline is to provide clinicians with information and treatment guidelines for sepsis in pregnancy. It particularly focuses on recognition of risk factors, diagnosis and management of sepsis in the peri partum period, including Early Goal Directed Therapy in the case of severe sepsis.


ISBN number:
Endorsed by:
Last Revised:
Contact:

978-1-74243-502-2
South Australian Maternal, Neonatal & Gynaecology Community of Practice
06/03/2017
HealthCYWHSPerinatalProtocol@sa.gov.au




Sepsis in Pregnancy

Flowchart: Initial management of maternal sepsis¹



Sepsis in Pregnancy

Flowchart: Early goal directed therapy for sepsis during pregnancy²



Sepsis in Pregnancy

EGDT (early goal directed therapy) - goals and normal values in pregnancy chart³

Measures	Resuscitation goals	Normal third trimester physiologic values
Central venous pressure	8-12 mm Hg	4-10 mm Hg
Mean arterial pressure	65 mm Hg or more	84-96 mm Hg
Urine output	More than 0.5 mL / kg / hour	Minimum 0.5 mL / kg / hour
Mixed venous oxygen saturation	More than 70 %	More than 80 %
Heart rate	Decreasing in response to treatment	83 (+/- 10) beats / minute

Sepsis in Pregnancy

Table of Contents

[Flowchart: Initial Management of Maternal Sepsis](#)

[Flowchart: Early Goal Directed Therapy for Sepsis during Pregnancy](#)

[EGDT goals and normal values in pregnancy chart](#)

[Summary of Practice Recommendations](#)

[Abbreviations](#)

[Definitions](#)

[Introduction](#)

[Common causes of sepsis during pregnancy](#)

[Common sites of infection in pregnancy](#)

[Other causes](#)

[Pathogens](#)

[Common bacterial infections and pathogens related to septic shock \(Table\)](#)

[Risk factors for maternal sepsis](#)

[Diagnosis of infection](#)

[Symptoms](#)

[Signs](#)

[Assessment for end-organ dysfunction \(Table\)](#)

[Evaluate potential sources of infection](#)

[Investigations](#)

[Blood tests](#)

[Microbiology](#)

[Imaging](#)

[Site of infection – specific considerations](#)

[Chorioamnionitis](#)

[Endometritis](#)

[Urinary tract infection](#)

[Mastitis](#)

[Wound, skin and soft tissue infections](#)

[Necrotising fasciitis](#)

[Bacteraemia and septic shock](#)

[Pneumonia](#)

[Gastroenteritis](#)

[Pharyngitis](#)

[Infection post termination of pregnancy /miscarriage](#)

[Cholecystitis / Cholangitis](#)

[Epidural abscess](#)

[Management](#)

[Surviving Sepsis Campaign Resuscitation Bundle \(Table\)](#)

[Initial management \(0-6 hours\)](#)

[Airway](#)

[Breathing](#)

[Circulation](#)

[Documentation and differential diagnosis](#)

ISBN number:

Endorsed by:

Last Revised:

Contact:

978-1-74243-502-2

South Australian Maternal, Neonatal & Gynaecology Community of Practice

06/03/2017

HealthCYWHSPerinatalProtocol@sa.gov.au

[Fetal wellbeing](#)

Sepsis in Pregnancy

[Remove the source of the maternal infection](#)

[Maternal observations](#)

[Antibiotics](#)

[Thromboprophylaxis](#)

[Staphylococcal and streptococcal toxic shock syndrome \(Table\)](#)

[Intravenous immunoglobulin \(IVIG\)](#)

[Complications](#)

[References](#)

[Useful online references](#)

[Acknowledgements](#)

Summary of Practice Recommendations

- > Sepsis may arise from many sources and is not limited to infections arising from the genital tract.
- > The onset of sepsis may be insidious; women with severe sepsis may appear very well before suddenly collapsing, with little or no warning.
- > Abdominal pain, fever ($> 38^{\circ}\text{C}$), tachypnoea and sustained tachycardia (> 90 beats per minute) are indications for admission and intravenous antibiotics
- > Diarrhoea and/or vomiting in pregnant women may be serious signs of sepsis and an indication for commencing antibiotic therapy.
- > Early investigation of non-specific symptoms in pregnant women is necessary to exclude serious infection.
- > Urinary tract infection and chorioamnionitis are common infections associated with septic shock in the pregnant woman.
- > Signs of severe sepsis in peripartum women, particularly with confirmed or suspected Group A streptococcal infection, should be regarded as an obstetric emergency
- > Early involvement of senior obstetrician / intensivist / anaesthetist / physician / microbiologist / infectious diseases consultant (as required) with appropriate referral and retrieval is essential. Suspicion of severe sepsis or septic shock should trigger urgent referral to a tertiary centre with adult intensive care facilities.
- > Follow the Surviving Sepsis Campaign Resuscitation 'Bundle' in severe sepsis.
- > Monitor and maintain airway, breathing and circulation as first priority.
- > In the event of hypotension and / or lactate > 4 mmol/L initiate urgent fluid resuscitation
- > Obtain blood cultures before antibiotic administration if possible but do not delay commencement of antibiotics
- > Systemically ill patients should commence broad spectrum intravenous antibiotics within 1 hour of recognition of severe sepsis
- > In a critically ill pregnant woman, birth of the baby may be considered if it would be beneficial to the mother or baby.

Sepsis in Pregnancy

Abbreviations

ARDS	Acute respiratory distress syndrome
AVPU	Alert, verbal, pain, unresponsive
SaO ₂	Arterial oxygen saturation
ASAP	As soon as possible
BP	Blood pressure
BSLs	Blood sugar levels
C	Celsius
CBP	Complete blood picture
CNS	Central nervous system
CVP	Central venous pressure
CVS	Cerebro-vascular system
EGDT	Early goal directed therapy
ECG	Electrocardiograph
GAS	Group A beta haemolytic streptococci
GCS	Glasgow coma scale
HCO ₃	Bicarbonate ion
HCT	Haematocrit
HVS	High vaginal swab
ICU	Intensive care unit
kg	Kilogram(s)
LFTs	Liver function tests
LVS	Low vaginal swab
MAP	Mean arterial blood pressure
mmHg	Millimetres of mercury
mL	Millilitre(s)
MSSU	Micro specimen of urine
PCO ₂	Carbon dioxide partial pressure
PO ₂	Oxygen partial pressure
PPROM	Preterm prelabour rupture of the membranes
RBC	Red blood cells
RCOG	Royal College of Obstetricians and Gynaecologists
ScvO ₂	Central venous oxygen saturation
SpO ₂	Oxygen saturation measured by pulse oximetry
SBP	Systolic blood pressure
SIRS	Systemic inflammatory response syndrome
WCC	White cell count
WHO	World Health Organisation

Sepsis in Pregnancy

Definitions

Systemic Inflammatory Response Syndrome (SIRS)	More than one of the following clinical findings: <ul style="list-style-type: none"> ◦ Temperature > 38°C or < 36°C ◦ Heart rate > 90 per minute ◦ Hyperventilation (evidence by respiratory rate > 20 per minute or PCO₂ < 32 mm Hg) ◦ WCC >12,000 or < 3,000
Sepsis	◦ Presence of both infection (invasion of tissue, fluid or a body cavity by pathogenic micro-organisms) and systemic manifestations of inflammatory response syndrome (SIRS)
Severe sepsis	◦ Sepsis complicated by sepsis-induced organ dysfunction or developing tissue hypoperfusion
Septic shock	<ul style="list-style-type: none"> ◦ Persistence of hypoperfusion (hypotension) in a septic patient, despite adequate volume resuscitation ◦ Hypotension is defined as: Systolic blood pressure (SBP) < 90 mm Hg or mean arterial blood pressure (MAP) < 60 mm Hg, or reduction of SBP > 40 mm Hg from baseline
Puerperal Sepsis	<ul style="list-style-type: none"> ◦ The term is used to describe sepsis occurring after delivery and the World Health Organisation (WHO)⁹ has defined it as “infection of the genital tract, occurring at any time between rupture of membranes or labour, and the 42nd day postpartum”, in which two or more of the following are present: <ul style="list-style-type: none"> > pelvic pain > fever > abnormal vaginal discharge > abnormal smell of discharge > delay in postpartum reduction of size of uterus

Introduction

- > Sepsis may arise in pregnancy at any time: before delivery, during labour or postpartum.
- > Sepsis may arise from many sources and is not limited to infections arising from the genital tract
- > The onset of sepsis may be insidious; women with severe sepsis may appear very well before suddenly collapsing, with little or no warning
- > Early identification of severe sepsis allows prompt, appropriate multidisciplinary management
- > Suspicion of severe sepsis or septic shock should trigger urgent referral to a tertiary centre with adult intensive care facilities
- > Up to 20 to 30 % of intensive care unit (ICU) admissions of obstetric patients result from sepsis in pregnancy¹¹
- > The contribution of sepsis as a cause of maternal mortality is between 3 % in developed countries and 12 % in developing countries¹¹
- > One third of early maternal mortality is due to refractory hypotension. Late maternal mortality is due to multiple organ failure¹¹
- > Urinary tract infection and chorioamnionitis are common infections associated with septic shock in the pregnant woman⁷
- > In the United Kingdom, the rapid course coupled with late presentation of some women with sepsis have contributed to a recent increase in direct maternal mortality due to genital tract sepsis (especially secondary to Group A streptococcal disease)⁵

Sepsis in Pregnancy

Common causes of sepsis in pregnancy

Common sites of infection in pregnancy

- > Urinary tract (pyelonephritis)
- > Pelvic structures (chorioamnionitis and endometritis)
- > Surgical wounds (caesarean section, perineal laceration)
- > Breast (mastitis)

Other causes

- > Infection of intravenous cannula sites
- > After urological procedures in the presence of urinary tract infection
- > Related to regional anaesthesia e.g. spinal / epidural abscess (rare)
- > Pneumonia (viral and bacterial)
- > Intrauterine fetal death
- > Septic abortion
- > Acute appendicitis
- > Acute cholecystitis
- > Pancreatitis
- > Bowel perforation (more common with inflammatory bowel disease)
- > Necrotising fasciitis
- > Meningitis

Common pathogens

- > The most prevalent bacterial organisms responsible for severe infection include:
 - > Group A beta haemolytic streptococci (GAS), also known as *Streptococcus pyogenes*
 - > Group B streptococcus
 - > *Escherichia coli*
 - > *Klebsiella*
 - > *Staphylococcus aureus*
 - > Anaerobes: peptostreptococci, peptococci, bacteroides, clostridium
- > *Clostridium* species and *listeria monocytogenes* are less common
- > Viral causes e.g. influenza, varicella, hepatitis and herpes simplex
- > Malaria and other tropical infections

Sepsis in Pregnancy

Common bacterial infections and pathogens related to septic shock¹⁰

Infections and usual pathogens (in brackets – see right column)	Pathogens
Pyelonephritis (1,4)	1. Escherichia coli
Pneumonia (6,7)	2. Bacteroides
Chorioamnionitis (1,2,8-12)	3. Clostridium
Endomyometritis (primarily after caesarean section) (1,2,5,9,12)	4. Klebsiella
Sepsis after miscarriage or termination of pregnancy (1,3)	5. Pseudomonas aeruginosa
Caesarean wound infection (1,2,6,7)	6. Streptococcus species
Severe mastitis (7)	7. Staphylococcus aureus
Necrotising fasciitis (2,3,6,9)	8. Group B streptococcus
	9. Peptostreptococcus
	10. Enterococcus
	11. Listeria monocytogenes
	12. Enterobacter

Risk factors for maternal sepsis^{13, 14}

- > Obesity
- > Impaired glucose tolerance / diabetes
- > Impaired immunity / immunosuppressant medication
- > Anaemia
- > Vaginal discharge
- > History of pelvic infection
- > History of group B streptococcal infection
- > Amniocentesis and other invasive procedures
- > Cervical Cerclage
- > Prolonged spontaneous rupture of membranes
- > Diabetes Mellitus
- > Group A streptococcal (GAS) infection in close contacts / family members (working with or having young children)
- > Of black or other minority ethnic group origin⁷
- > Retained products of conception
- > Caesarean birth (greater risk associated with emergency caesarean)

Diagnosis of infection

A thorough history and physical examination is required to make the diagnosis of sepsis. Important elements in making a diagnosis include:

1. Having a suspicion of the diagnosis of sepsis
2. Assessing for evidence of end organ dysfunction
3. Identifying site / source of the infection
4. The speed of onset or deterioration in symptoms and signs is important

Symptoms

- > Chills, sweating, warm skin, faintness or syncope, vomiting, rash, headache, dyspnoea, general weakness (or pain related to sites of sepsis – see below)

Sepsis in Pregnancy

Signs

One or more of the following:

- > Pyrexia (38°C or more): May not always be present and is not necessarily related to the severity of sepsis. Temperature may be less than 36°C in severe sepsis
- > Tachypnoea (20 breaths or more per minute)
- > Hypoxia
- > Tachycardia (90 or more bpm)
- > Arterial hypotension (systolic blood pressure less than 90 mm Hg; mean arterial pressure less than 70 mm Hg; or systolic blood pressure decrease 40 or more mm Hg)
- > Decreased capillary refill or mottling
- > Fetal distress secondary to maternal acidosis
- > Oliguria (urine output less than 0.5 mL / kg / hr for at least two hours, despite adequate fluid resuscitation)
- > Considerable oedema or positive fluid balance (20 or more mL / kg over 24 hours)
- > Ileus (absent bowel sounds)
- > Impaired mental state, altered conscious level
- > Hyperglycaemia in the absence of diabetes (plasma glucose more than 7.7 mmol / L)
- > Bruising or discolouration of skin suggests late fasciitis (often pain receding as cutaneous anaesthesia supervenes as nerves die)
- > Failure to respond to treatment

Assessment for end-organ dysfunction	
Acute circulatory failure	Cardiovascular assessment: Peripheral perfusion, pulse, BP, CVP including signs of high or low cardiac output
Metabolic acidosis	pH, HCO ₃ , PaCO ₂ , lactate, anion gap
Acute hypoxaemia	SpO ₂ , PaO ₂ , PaO ₂ / FIO ₂
Liver dysfunction	LFTs
Coagulopathy	CBP and coagulation profile
Acute renal failure	Fluid overload, urine output Urea, creatinine, electrolytes
CNS disturbance e.g. altered mental state	Sedation score AVPU GCS Encephalopathy
Overall metabolism	Assessment for a hyper-catabolic state: negative nitrogen balance (e.g. low albumin) or raised blood glucose levels

Sepsis in Pregnancy

Evaluate potential sources of infection

- > Identify if any recent sore throat or respiratory illness or close contact with persons with illness (particularly streptococcal infections)
- > Exclude intravenous drug misuse (high risk of staphylococcal and streptococcal sepsis)
- > Recent febrile illness (especially if associated with chills and rigors -suggest bacteraemia or viraemia)
- > Gastrointestinal symptoms (e.g. diarrhoea and vomiting) may be due to food borne pathogens or early toxic shock
- > Consider need for vaginal examination to exclude retained tampons or swabs
- > Exclude multi-resistant organisms such as ESBL-producing gram negative bacteria, vancomycin-resistant enterococci and MRSA (may affect choice of antimicrobial to combat sepsis and infection control precautions)
- > Early investigation of non-specific symptoms in pregnant women is necessary to exclude serious infection.¹⁷

Investigations

Blood tests	
Serum lactate	Greater than 4 mmol/L requires urgent action
Complete blood picture	White blood cell (WBC) count $> 12 \times 10^9$ Leucopenia - WBC count $< 4 \times 10^9$ Normal WBC count with $> 10\%$ immature forms
Plasma C-reactive protein	More than 7 mg/L (usually significantly higher in bacterial sepsis)
Urea and electrolytes	Creatinine rise of > 44.2 micromol/L; sepsis is severe if creatinine level > 176 micromol/L
Plasma glucose	Hyperglycaemia in the absence of diabetes (plasma glucose > 7.7 mmol/L)
Liver function tests (LFTs)	Hyperbilirubinaemia (plasma total bilirubin > 70 micromol/L)
Coagulation profile	Coagulation abnormalities (INR > 1.5 or APTT > 60 seconds)
Blood gas Particularly for respiratory tract sepsis but also may detect acidosis from early stage shock	Arterial hypoxaemia ($\text{PaO}_2 / \text{FIO}_2 < 300$ mm Hg) Sepsis is severe if < 250 mm Hg in the absence of pneumonia or < 200 mm Hg in the presence of pneumonia Raised serum lactate ≥ 4 mmol/L

Adapted from: RCOG Diagnostic criteria for sepsis (modified from Levy⁶)

Sepsis in Pregnancy

Microbiology

- > Blood cultures (ideally obtain before antibiotic administration)
- > Obtain cultures of additional sites as indicated and as soon as possible e.g. midstream specimen urine (MSSU), wound swab (episiotomy or caesarean section), placental swabs, respiratory secretions, naso-pharyngeal aspirate (NPA), amniotic fluid, cerebrospinal fluid, HVS, LVS, endocervical, expressed breast milk
- > If the methicillin-resistant *Staphylococcus aureus* (MRSA) status is unknown, obtain swabs from nose, groin and axilla and send for urgent screening
- > Suspected pneumonia - Chest X-ray, naso-pharyngeal aspirate

Imaging

- > Consider imaging modalities e.g. Chest X-ray, pelvic ultrasound, computed tomography, magnetic resonance imaging (may help define inflammation or the collection of pus)
- > Echocardiography may be useful in cases of women who are intravenous drug users, particularly those with staphylococcal bacteraemia as there may be right-sided endocarditis and to assess cardiac function

Site of infection – specific considerations

Early diagnosis to determine if surgical intervention required

May be difficult to localise the source

Chorioamnionitis

Diagnosis of chorioamnionitis relies on the clinical presentation

- > Maternal fever > 38° C with any 2 of the following:
 - > Increased white cell count (>15 x 10⁹/L)
 - > Maternal tachycardia (> 100 bpm)
 - > Fetal tachycardia (>160 bpm)
 - > Uterine tenderness
 - > Offensive smelling vaginal discharge
 - > C-Reactive Protein > 40
 - > Histological examination of placenta and membranes with evidence of acute inflammation may confirm diagnosis after birth

Treatment of chorioamnionitis

- > Amoxicillin 2 g IV every 6 hours
- > Gentamicin 5 mg/kg IV daily
- > Metronidazole 500 mg IV every 12 hours
- > If allergic to penicillin, give clindamycin 450 mg IV in 50 – 100 mL over at least 20 minutes every 8 hours AND gentamicin 5 mg / kg IV daily until delivery
- > Do not inhibit labour, but consider hastening delivery under intravenous antibiotic cover
- > Consider optimal mode of delivery (LSCS versus vaginal birth) on the basis of the findings and the anticipated duration until birth
- > Consider ongoing treatment with antibiotics postnatally

Endometritis

- > Offensive lochia
- > Unusual bleeding
- > Consider need for evacuation of retained products of conception
- > If does not respond to antibiotics, consider septic pelvic thrombosis and commence on heparin

Sepsis in Pregnancy

Urinary tract infection

- > Loin pain
- > Tenderness to renal percussion
- > Send midstream specimen of urine (MSSU)
- > Treat acute pyelonephritis aggressively (For more information see 'Urinary tract infections in pregnancy' in the A to Z index at www.sahealth.sa.gov.au/perinatal)

Mastitis

- > Reddened sector in the breast
- > Immediate referral to hospital if the woman is clinically unwell, no response to oral antibiotics within 48 hours, mastitis recurs or there are severe or unusual symptoms
- > Complications include breast abscess, necrotising fasciitis, toxic shock syndrome

Wound, skin and soft tissue infections

- > Signs of inflammation at the site
- > Fever, chills, tachycardia, tachypnoea, wound swelling, warm to touch, painful, discharge
- > Closed space infections need surgical drainage
- > Carefully examine intravenous cannulae or injection sites, and caesarean or episiotomy wounds
- > Take a swab of any discharge
- > Exclude complications of venous thrombosis or necrotising fasciitis (see below)

Necrotising fasciitis

- > Extreme pain, associated with signs of prostration and severe sepsis
- > Requires early surgical intervention with fasciotomy and aggressive antibiotic treatment

Bacteraemia and septic shock

- > Suspected bacterial sepsis requires careful examination for skin and soft tissue infection
- > If drains, vascular access devices or other indwelling devices are suspected as the source of infection, remove as soon as is practicable
- > Skin and soft tissue infections are particularly associated with toxic shock syndromes
- > Septicaemic seeding of streptococci from a uterine focus may give rise to a secondary focus in a limb, simulating a venous thrombosis
- > Women with thrombosis who are systemically unwell with any features of sepsis should be examined very carefully. If shock or other organ dysfunction present, rapid referral to adult intensive care

Pneumonia

- > Fever, chills, dyspnoea, cough, sputum production, pleuritic pain
- > Consider referral to respiratory physician, medical microbiologist
- > Send sputum sample, throat swab for culture
- > Chest X-ray

Gastroenteritis

- > Diarrhoea and vomiting may be symptoms of bacterial sepsis, due to organisms including staphylococcus and streptococcus
- > Stool sample for routine culture (include C. difficile toxin testing if diarrhoea offensive after antimicrobial treatment)
- > Salmonella and Campylobacter rarely cause severe systemic infection

Sepsis in Pregnancy

Pharyngitis

- > Usually viral
- > Approximately 10 % of cases are due to GAS
- > Obtain throat swab for culture
- > Antibiotic treatment if 3 or more of the following: fever, tonsillar exudate, no cough, tender anterior cervical lymphadenopathy

Infection post termination of pregnancy / miscarriage

- > Suspect infection in women with a history of a recent termination of pregnancy or spontaneous miscarriage who present with pyrexia, persistent bleeding or abdominal pain, especially if the pain is constant and severe
- > Vaginal swabs, ultrasound scan to exclude retained products of conception and consider diagnostic evacuation of uterus (evacuation of retained products of conception) if there is still doubt
- > Complete blood picture, C-reactive protein, blood cultures if pyrexia > 38° Celsius
- > Commence high dose broad-spectrum intravenous antibiotics without waiting for microbiology results

Cholecystitis / Cholangitis

- > Upper right quadrant pain, biliary colic after a fatty meal, jaundice, nausea, vomiting, fever, ultrasonography examination (USG) of abdomen

Epidural abscess

- > Very rare complication
- > *S. aureus* is the usual organism
- > Severe backache
- > Neurological deficit including bowel / bladder dysfunction and unexplained fever

Management

- > After identification of severe sepsis, in the first 6 hours, follow the Surviving Sepsis Campaign Resuscitation 'Bundle' (see below)
- > Treat underlying infection and support failing organ functions
- > Systemically ill patients should commence broad spectrum, intravenous antibiotics (within 1 hour of recognition of severe sepsis)
- > Patients with single organ failure not responding to simple measures require transfer to a tertiary centre with high dependency care
- > Patients with two or more organ failures and respiratory failure require transfer to a tertiary centre with Adult Intensive Care Facilities (see 'Assess for evidence of end-organ dysfunction' above)
- > **Key Practice Point for outpatients:**
Abdominal pain, fever (> 38°C), tachypnoea and sustained tachycardia (> 90 beats per minute) are indications for admission and intravenous antibiotics

Sepsis in Pregnancy

Surviving Sepsis Campaign Resuscitation 'Bundle'

Tasks to be performed within the first 6 hours of the identification of severe sepsis⁷

- > Obtain blood cultures before antibiotic administration
- > Administer broad spectrum antibiotic within one hour of recognition of severe sepsis
- > Measure serum lactate
- > In the event of hypotension and / or lactate > 4 mmol/L:
 - > Deliver an initial minimum 20 mL / kg of crystalloid or an equivalent fluid
 - > Apply vasopressors for hypotension not responding to initial fluid resuscitation to maintain mean arterial pressure above 65 mm Hg
- > In the event of persistent hypotension despite fluid resuscitation (septic shock) and / or serum lactate > 4 mmol/L:
 - > Achieve a central venous pressure (CVP) of ≥ 8 mm Hg
 - > Achieve a central venous oxygen saturation (ScvO₂) ≥ 70 % or mixed venous oxygen saturation (ScvO₂) ≥ 65 %

Adapted from the Surviving Sepsis Campaign Resuscitation Bundle (group of therapies)¹⁵

Initial management (0-6 hours)

- > Early involvement of senior obstetrician / intensivist / anaesthetist / physician / microbiologist / infectious diseases consultant (as required) with appropriate consultation, referral and retrieval
- > Assess need for central venous access and initiate Surviving Sepsis Campaign Resuscitation Bundle if severe sepsis is suspected (see table 3 above)
- > Follow the A, B, C, D, E's of resuscitation

Airway

- > Maintain a clear airway
- > Administer oxygen via non-rebreathing mask at 12 litres per minute
- > If the woman has collapsed, check her airway is patent, she is breathing and maintain her position in left-lateral

Breathing

- > Assess breathing pattern, rate and colour and ventilate if required
- > Attach pulse oximeter, electrocardiogram and automatic blood pressure monitors and monitor SpO₂, maternal pulse and blood pressure

Circulation

- > Hypovolaemia is present in almost all patients with septic shock
- > Obtain intravenous access
- > Take blood for blood cultures, serum lactate, complete blood count, coagulation profile, urea and electrolytes, liver function tests, blood glucose
- > Immediate fluid resuscitation with either crystalloid or colloid as per medical order
- > If no response to simple measures of fluid resuscitation consider the need for insertion of central venous catheter and CVP monitoring
- > **Correct hypotension** – Administer vasopressors for hypotension that is not responding to initial fluid resuscitation to maintain mean arterial pressure (MAP) more than 65 mm Hg (intensivist or anaesthetic management)
- > Perform arterial blood gases

Sepsis in Pregnancy

Documentation and differential diagnosis

- > Obtain history as soon as possible
- > Document vital signs on 'Rapid deterioration and Response' chart
- > Confirm diagnosis
- > Commence broad spectrum intravenous antibiotic treatment within 1 hour of diagnosis of severe sepsis
- > Commence fluid balance chart and monitor urine output

NB: Fluid overload may lead to fatal pulmonary or cerebral oedema in women with septic shock. Clear, accurate documentation and careful monitoring of fluid balance is essential

Fetal wellbeing

- > Assess and monitor fetal wellbeing (cardiotocography) if applicable
- > Continuous fetal monitoring intrapartum
- > Consider the need for delivery as applicable
 - > In a critically ill pregnant woman, birth of the baby may be considered if it would be beneficial to the mother or the baby or to both. A decision on the timing and mode of birth should be made by a senior obstetrician after discussion with the woman if her condition allows
 - > If preterm birth is anticipated, cautious consideration should be given to the use of antenatal corticosteroids for fetal lung maturity in the woman with sepsis
 - > Epidural / spinal anaesthesia should be avoided in women with sepsis

Remove the source of the maternal infection

- > Expedite delivery if there are signs of chorioamnionitis
- > Any retained products of conception should be removed as soon as the maternal condition is stable
- > If large amounts of pelvic pus are present a laparotomy and sometimes hysterectomy may be necessary

Maternal observations

- > Obtain and document temperature, pulse, respiratory rate, blood pressure, SpO₂ every 15 minutes in the initial assessment phase
 - > Once stable, continue hourly
- > Utilise 'Rapid Deterioration and Response' chart. The use of early warning charts and escalation guidelines including involvement of senior medical staff (intensivist, physician) assists in the early detection and management of the deteriorating patient
- > Glasgow Coma Scale and pupil response if required
- > Hourly urine output (consider insertion of indwelling catheter)
- > In rapidly deteriorating cases, ensure urgent referral to critical care team and obstetric consultant

Sepsis in Pregnancy

Antibiotics

- > Early consult with microbiologist or infectious diseases consultant for women with evidence of systemic infection
- > Commence broad spectrum intravenous antibiotic cover within 1 hour of suspicion of severe sepsis, with or without septic shock
- > The choice of antibiotic depends on the clinical suspicion, local flora and culture information, if available
- > If genital tract sepsis is suspected, prompt early treatment with a combination of high-dose broad-spectrum intravenous antibiotics may be lifesaving
- > Empirical treatment should include broad spectrum antimicrobials active against Gram-negative bacteria, and capable of preventing exotoxin production from Gram-positive bacteria (according to local microbiology policy). Gram-positive cover is necessary if the likelihood of this infection is high

Thromboprophylaxis

- > Women with sepsis are at increased risk of venous thromboembolism
- > Consider prophylaxis with low molecular weight heparin
- > Use of compression stockings

Staphylococcal and streptococcal toxic shock syndrome

Toxic shock syndrome is a life-threatening illness characterised by hypotension and multi-organ failure. It may be caused by *Staphylococcus aureus* (rarely isolated) or *Streptococcus pyogenes* that produce and release superantigenic exotoxins. The exotoxins activate T-cells on a large scale resulting in a massive release of inflammatory cytokines¹⁶

Staphylococcal toxic shock syndrome	Streptococcal toxic shock syndrome
<ol style="list-style-type: none"> 1. Fever $\geq 39.9^{\circ}\text{C}$ 2. Rash: diffuse macular erythema 3. Desquamation: 10-14 days after onset of illness, especially palms and soles 4. Hypotension: systolic BP < 90 mmHg 5. Multisystem involvement: 3 or more of the following systems affected: <ul style="list-style-type: none"> o GIT: vomiting or diarrhoea at onset of illness o Muscular: severe myalgia or elevated creatinine phosphokinase o Mucous membranes: vaginal, oropharyngeal or conjunctival hyperaemia o Renal: creatinine twice the upper limit of normal o Haematological – platelets $\leq 100 \times 10^9/\text{L}$ o Central nervous system – disorientation or alterations in consciousness without focal neurological signs 	<p>A. Isolation of group A Streptococcus from:</p> <ol style="list-style-type: none"> 1. Normally sterile site: blood, cerebrospinal fluid, peritoneal fluid, tissue biopsy 2. Non-sterile site: throat, vagina, sputum <p>B. Clinical case definition</p> <p>Multi-organ involvement characterised by:</p> <ol style="list-style-type: none"> 1. Hypotension and 2. 2 or more of the following: <ul style="list-style-type: none"> o Renal impairment – creatinine > 176 micromol/L o Coagulopathy – platelets $< 100 \times 10^9/\text{L}$ or disseminated intravascular coagulation o Liver involvement: alanine transaminase or aspartate transaminase or bilirubin levels twice the normal upper limit for age o Acute respiratory distress syndrome o Generalised erythematous macular rash (present in 10 %): may desquamate o Soft tissue necrosis including necrotising fasciitis, myositis or gangrene
Case classification:	Case classification
<ul style="list-style-type: none"> o Probable: 4 of the 5 clinical findings positive o Confirmed: case with all 5 clinical findings 	<ul style="list-style-type: none"> o Probable: meets clinical case definition (above) plus isolation from non-sterile site o Definite: meets clinical case definition (above) plus isolation of group A Streptococcus from a normally sterile site

Sepsis in Pregnancy

Intravenous immunoglobulin (IVIG)

- > IVIG is recommended for severe invasive streptococcal or staphylococcal infection if other therapies have failed
- > High dose IVIG has been used in pregnant and postpartum women and is effective in exotoxic shock (i.e. toxic shock attributable to streptococci and staphylococci)⁸
- > For more information, see 'Intragam infusion' in the A to Z index at www.sahealth.sa.gov.au/perinatal

Complications

- > Acute respiratory distress syndrome (ARDS)
- > Disseminated Intravascular coagulation
- > Renal failure
- > Hepatic failure

Sepsis in Pregnancy

References

1. Winter C, Crofts J, Laxton C, Barnfield S, Draycott T, United Kingdom editors. Practical Obstetric Multi-Professional Training (PROMPT) course manual. Practical locally based training for obstetric emergencies. Australian and New Zealand edition. Sowter M, Weaver E, Beaves M, Australian editors. Melbourne: RANZCOG; 2013.
2. Otero RM, Nguyen HB, Huang DT, Gaieski DF, Goyal M, Gunnerson KJ, et al. Early goal-directed therapy in severe sepsis and septic shock revisited concepts, controversies, and contemporary findings. *Chest* 2006; 130: 1579-95
3. Norwitz ER, Robinson JN, Malone FD. Pregnancy-induced physiologic alterations. In: Dildy GA III, Belfort MA, Saade G, et al, editors. *Critical care obstetrics*. 4th edition. Malden: Blackwell Science; 2004. p. 19-42
4. Lucas DN, Robinson PN, Nel MR. Sepsis in obstetrics and the role of the anaesthetist. *Int J Obstet Anesthesia* 2012; 21: 56-67.
5. Harper A. Chapter 7: Sepsis. In: Centre for Maternal and Child Enquiries (CMACE). *Saving Mother's Lives: reviewing maternal deaths to make motherhood safer: 2006-2008*. *BJOG* 2011; 118 (suppl. 1): 85-96.
6. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003; 31: 1250-6. (Consensus)
7. Royal College of Obstetricians and Gynaecologists (RCOG). Bacterial sepsis in pregnancy – 64a April 2012. RCOG Green top Guidelines. London: RCOG Press; 2012a. Available from URL: <http://www.rcog.org.uk/womens-health/clinical-guidance/sepsis-pregnancy-bacterial-green-top-64a>
8. Royal College of Obstetricians and Gynaecologists (RCOG). Bacterial sepsis following pregnancy – 64b April 2012. RCOG Green top Guidelines. London: RCOG Press; 2012b. Available from URL: <http://www.rcog.org.uk/womens-health/clinical-guidance/sepsis-following-pregnancy-bacterial-green-top-64b>
9. Dolea C, Stein C. Global burden of maternal sepsis in the year 2000. Evidence and information for policy. World Health Organisation (WHO), Geneva; July 2003.
10. Joseph J, Sinha A, Paech M, Walters BNJ. Sepsis in pregnancy and early goal-directed therapy. *Obstet medicine* 2009; 2: 93-9.
11. Prasad BGR, Sunanda GV. Sepsis. In: Johanson R, Cox C, Grady K, Howell C, editors. *Managing obstetric emergencies and trauma. The MOET Course Manual*. London: RCOG Press; 2003. p. 231-234.
12. Funk D, Sebat F, Kumar A. A systems approach to the early recognition and rapid administration of best practice therapy in sepsis and septic shock. *Current Opinion in Critical Care* 2009; 15: 301-07.
13. Lewis G, editor. *Saving Mothers' lives: Reviewing maternal deaths to make motherhood safer – 2003-2005. The Seventh Report on Confidential enquiries into maternal deaths in the United Kingdom*. London: RCOG Press; 2007.
14. Centre for Maternal and Child Enquiries. CMACE Emergent theme briefing #1: Genital tract sepsis. *Saving Mothers' Lives 2006-08: Briefing on genital tract sepsis*. London: CMACE; 2010.
15. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R et al. Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock. *Crit Care Med* 2008; 36: 296-327
16. National Blood Authority (NBA). Jurisdictional Blood Committee, for and on behalf of the Health Minister's Conference. Criteria for the clinical use of intravenous immunoglobulin in Australia, 2nd edition. Canberra: Commonwealth of Australia; 2012. Available from URL: <http://www.blood.gov.au/system/files/documents/nba-ivig-criteria-for-use-2nd-edition.pdf>
17. AIHW: Humphrey MD, Bonello MR, Chughtai A, Macaldowie A, Harris K & Chambers GM 2015. Maternal deaths in Australia 2008–2012. Maternal deaths series no. 5. Cat. no. PER 70. Canberra: AIHW.

Sepsis in Pregnancy

Useful online references

Royal College of Obstetricians and Gynaecologists (RCOG). Bacterial sepsis in pregnancy – 64a April 2012. RCOG Green top Guideline. Available from URL:

<http://www.rcog.org.uk/womens-health/clinical-guidance/sepsis-pregnancy-bacterial-green-top-64a>

College of Intensive Care Medicine of Australia and New Zealand: IC-01 Minimum standards for Intensive Care 2011. Available from URL:

http://www.cicm.org.au/CICM_Media/CICMSite/CICM-website/Resources/Professional%20Documents/IC-1-Minimum-Standards-for-Intensive-Care-Units.pdf

Surviving Sepsis Campaign at <http://www.survivingsepsis.org>

Sepsis in Pregnancy

Acknowledgements

The Royal College of Obstetricians and Gynaecologists Green top Guideline 64a - Bacterial sepsis in pregnancy (April 2012) has largely been used to inform this guideline

The South Australian Perinatal Practice Guidelines gratefully acknowledge the contribution of clinicians and other stakeholders who participated throughout the guideline development process particularly:

Write Group Leads

Prof Marc Keirse
Dr Kym Osborn
Allison Rogers
Dr Steven Scroggs

Write Group Members

SAPPG Management Group
Dr Ranjan Joshi

SAPPG Management Group Members

Sonia Angus
Dr Kris Bascomb
Lyn Bastian
Dr Feisal Chenia
John Coomblas
A/Prof Rosalie Grivell
Dr Sue Kennedy-Andrews
Jackie Kitschke
Catherine Leggett
Dr Anupam Parange
Dr Andrew McPhee
Rebecca Smith
Dr Nigel Stewart
Simone Stewart-Noble
A/Prof John Svigos
Dr Laura Willington

Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	20 Nov 2012	28 Feb 17	Original version
2.0	28 Feb 17	6 March 2017	Reviewed
3.0	6 March 2017	Current	Review and update